

Calculation of, and bounds for, the multipole moments of stationary spacetimes

Thomas Bäckdahl*, Magnus Herberthson*

Abstract

In this paper the multipole moments of stationary asymptotically flat spacetimes are considered. We show how the tensorial recursion of Geroch and Hansen can be replaced by a scalar recursion on $\mathbb{R}^2$. We also give a bound on the multipole moments. This gives a proof of the "necessary part" of a long standing conjecture due to Geroch.

1 Introduction

The relativistic multipole moments of asymptotically flat spacetimes have been defined by Geroch [6] for static spacetimes and then generalised to the stationary case by Hansen [7]. Together with Beig's [1] generalised definition of centre of mass this gives a coordinate independent description of all asymptotically flat stationary spacetimes.

However, the tensorial recursion which defines the multipole moments (1), is computationally rather complicated as it stands. In the axisymmetric (static or stationary) case, on the other hand, it was shown, [3], [4] that the recursion can be replaced by a scalar recursion on $\mathbb{R}$, and that all the moments can be collected into one complex valued function y on $\mathbb{R}$, where the moments are given by the the derivatives of y at 0.

In the general case, the multipole of order 2^n has $2n + 1$ degrees of freedom, as compared to one degree of freedom in the axisymmetric case. Therefore, apart from the technical problems, it is not obvious what form a generalisation to the general case should take. In this paper, we show that also in the general stationary case, the recursion (1) can be simplified to a scalar recursion, this time on $\mathbb{R}^2$. This is shown using normal coordinates, complex null geodesics, and exploiting the extra conformal freedom of the conformal compactification.

Using this simplification we can partially confirm an extension of a long standing conjecture by Geroch [6]:

Given any set of multipole moments, subject to the appropriate convergence condition, there exists a static solution of Einstein's equations having precisely those moments.

This conjecture has its natural extension to the stationary case.

*Department of Mathematics, Linköping University, SE-581 83 Linköping, Sweden.
e-mail: thbac@mai.liu.se, maher@mai.liu.se

In this paper we will state the appropriate convergence condition in the general stationary case, i.e., we will prove that this condition is necessary for existence of a stationary solution to Einstein's equations.

2 Multipole moments of stationary spacetimes

In this section we quote the definition of multipole moments given by Hansen in [7], which is an extension to stationary spacetimes of the definition by Geroch [6]. We thus consider a stationary spacetime (M, g_{ab}) with timelike Killing vector field ξ^a . We let $\lambda = -\xi^a \xi_a$ be the norm, and define the twist ω through $\nabla_a \omega = \epsilon_{abcd} \xi^b \nabla^c \xi^d$. If V is the 3-manifold of trajectories, the metric g_{ab} (with signature $(-, +, +, +)$) induces the positive definite metric

$$h_{ab} = \lambda g_{ab} + \xi_a \xi_b$$

on V . It is required that V is asymptotically flat, i.e., there exists a 3-manifold $\hat{V}$ and a conformal factor Ω satisfying

- (i) $\hat{V} = V \cup \Lambda$, where Λ is a single point
- (ii) $\hat{h}_{ab} = \Omega^2 h_{ab}$ is a smooth metric on $\hat{V}$
- (iii) At Λ , $\Omega = 0$, $\hat{D}_a \Omega = 0$, $\hat{D}_a \hat{D}_b \Omega = 2\hat{h}_{ab}$,

where $\hat{D}_a$ is the derivative operator associated with $\hat{h}_{ab}$. On M , and/or V one defines the scalar potential

$$\phi = \phi_M + i\phi_J, \quad \phi_M = \frac{\lambda^2 + \omega^2 - 1}{4\lambda}, \quad \phi_J = \frac{\omega}{2\lambda}.$$

The multipole moments of M are then defined on $\hat{V}$ as certain derivatives of the scalar potential $\hat{\phi} = \phi/\sqrt{\Omega}$ at Λ . More explicitly, following [7], let $\hat{R}_{ab}$ denote the Ricci tensor of $\hat{V}$, and let $P = \hat{\phi}$. Define the sequence $P, P_{a_1}, P_{a_1 a_2}, \dots$ of tensors recursively:

$$P_{a_1 \dots a_n} = C[\hat{D}_{a_1} P_{a_2 \dots a_n} - \frac{(n-1)(2n-3)}{2} \hat{R}_{a_1 a_2} P_{a_3 \dots a_n}], \quad (1)$$

where $C[\cdot]$ stands for taking the totally symmetric and trace-free part. The multipole moments of M are then defined as the tensors $P_{a_1 \dots a_n}$ at Λ . The requirement that all $P_{a_1 \dots a_n}$ be totally symmetric and trace-free makes the actual calculations very cumbersome.

In [2], [9] it was shown that (when the mass is non-zero) there exist a conformal factor Ω and a chart, such that all components of the metric $\hat{h}_{ab}$ and the potential $\hat{\phi}$ are analytic in terms of the coordinates, in a neighbourhood of the infinity-point. Expressed in these coordinates, the exponential map becomes analytic. Therefore, we can use Riemannian normal coordinates and still have analyticity of the metric components and the potential. If the mass is zero, this analyticity condition will be assumed. Thus, henceforth we assume that Ω is chosen such that the (rescaled) metric and potential are analytic in a neighbourhood of Λ .

3 Multipole moments through a scalar recursion on $\mathbb{R}^2$

Suppose that $(x^1, x^2, x^3) = (x, y, z)$ are normal coordinates (with respect to $\hat{h}_{ab}$) centred around Λ . This means that for any constants $a = a^1, b = a^2, c = a^3$ the curve $t \rightarrow (at, bt, ct)$ is a geodesic, i.e. in terms of coordinates that

$$\ddot{x}^i + \Gamma_{kl}^i \dot{x}^k \dot{x}^l = \Gamma_{kl}^i a^k a^l = 0$$

where the Christoffel symbols are evaluated at (at, bt, ct) for appropriate t . Due to analyticity, this relation holds for complex values of a, b, c , i.e., we can consider geodesics in the complexification $\hat{V}_{\mathbb{C}}$ of $\hat{V}$. Of particular interest is the one-parameter family of curves:

$$\gamma_\varphi : t \rightarrow (t \cos \varphi, t \sin \varphi, it), \quad t \in [0, t_0], \varphi \in [0, 2\pi)$$

for some suitable t_0 . The tangent vector $\eta^a = \eta_\varphi^a(t) = \cos \varphi (\frac{\partial}{\partial x^1})^a + \sin \varphi (\frac{\partial}{\partial x^2})^a + i(\frac{\partial}{\partial x^3})^a$ is seen to be a complex null vector along γ_φ . Namely, from $\eta^a \hat{D}_a \eta^b = 0$, we infer that $\eta^a \hat{D}_a (\eta^b \eta_b) = 0$. The (constant) value of $\eta^b \eta_b$ is then found to be 0 by evaluation at $t = 0$.

Next, consider the mapping $F : \mathbb{R}^2 \rightarrow V_{\mathbb{C}} : (\xi, \zeta) \rightarrow (\xi, \zeta, i\sqrt{\xi^2 + \zeta^2})$. We let S denote the 2-surface $F(U) \subset \hat{V}_{\mathbb{C}}$, where $U \subset \mathbb{R}^2$ is a suitable neighbourhood of $(\xi, \zeta) = (0, 0)$. S is then a smooth surface, except at Λ where it has a vertex point, closely resembling a null cone in a three dimensional Lorentzian space. The curves γ_φ are given by $\gamma_\varphi(t) = F(t \cos \varphi, t \sin \varphi)$, and in particular η^a lies along S . This suggests that we use the polar coordinates ρ, φ around Λ on S defined via $\xi = \rho \cos \varphi, \zeta = \rho \sin \varphi$. We now follow the approach from [4], where a useful vector field η^a on $\hat{V}$ was introduced. In [4], where the spacetime was axisymmetric, η^a was explicitly expressed in terms of the metric cast in the Weyl-Papapetrou form [10], and was defined on the whole of $\hat{V}$ except on the symmetry axis. In this paper the axisymmetry condition is dropped, which makes the construction of a corresponding η^a more difficult. In addition, a general spacetime has $2n + 1$ degrees of freedom for the multipole moment of order 2^n , compared to one degree of freedom in the axisymmetric case [8]. Nevertheless, it will turn out to be sufficient to know the potential $\hat{\phi}$ on S to determine all the moments. On S , η^a has the following properties.

Lemma 1. *Suppose $\hat{V}$ and S are defined as above. Then there exists a regularly direction dependent (at Λ) vector field η^a on S with the following properties:*

- a) $\eta^a \hat{D}_a \eta^b$ is parallel to η^b .
- b) For all tensors $T_{a_1 \dots a_n}$, $\eta^{a_1} \dots \eta^{a_n} T_{a_1 \dots a_n} = \eta^{a_1} \dots \eta^{a_n} C[T_{a_1 \dots a_n}]$,
- c) At Λ , $P_{a_1 \dots a_n}$ (in $\hat{V}$) is determined by $\eta^{a_1} \dots \eta^{a_n} P_{a_1 \dots a_n}$ (on S)

Proof. a) was demonstrated above, b) follows as in [4], while c) requires a different argument. A totally symmetric and trace-free tensor $P_{a_1 \dots a_n}$ has $2n + 1$ degrees of freedom, and in Cartesian coordinates (x, y, z) it can be expressed via the components $P_{\underbrace{x \dots x}_j \underbrace{y \dots y}_{n-j-1} z}$ and $P_{\underbrace{x \dots x}_j \underbrace{y \dots y}_{n-j}}$.¹ Therefore, at Λ , we

¹In brief, any index occurrence of several z 's can be removed via $P_{zz\dots} + P_{xx\dots} + P_{yy\dots} = 0$.

can write

$$\begin{aligned}
P_{a_1 \dots a_n} &= \sum_{j=0}^n a_j C[(dx)_{a_1} \dots (dx)_{a_j} (dy)_{a_{j+1}} \dots (dy)_{a_n}] \\
&+ \sum_{j=0}^{n-1} b_j C[(dx)_{a_1} \dots (dx)_{a_j} (dy)_{a_{j+1}} \dots (dy)_{a_{n-1}} (dz)_{a_n}]
\end{aligned} \tag{2}$$

Contracting with $\eta^{a_1} \dots \eta^{a_n}$, and using lemma 1b we find that

$$\eta^{a_1} \dots \eta^{a_n} P_{a_1 \dots a_n} = \sum_{j=0}^n a_j \cos^j \varphi \sin^{n-j} \varphi + i \sum_{j=0}^{n-1} b_j \cos^j \varphi \sin^{n-1-j} \varphi \tag{3}$$

If the left hand side is zero, the trigonometric polynomial to the right must be identically zero. This means that all the coefficients a_n and b_n are zero, and by (2) that $P_{a_1 \dots a_n}$ is zero. In particular the $2n+1$ components in the RHS of (2) are linearly independent. This proves c). $\square$

Note that although the moments are encoded in the coefficients a_n and b_n , this encoding is dependent on the choice of normal coordinates, i.e., the orientation of the coordinate axes in $T_\Lambda \hat{V}_\mathbb{C}$.

We can now replace the recursion (1) on $\hat{V}$ with a scalar recursion on S . Again, we follow [4], and define

$$f_n = \eta^{a_1} \eta^{a_2} \dots \eta^{a_n} P_{a_1 a_2 \dots a_n}, \quad n = 0, 1, 2, \dots \tag{4}$$

on S . In particular, $f_0 = P = \hat{\phi} = \phi/\sqrt{\Omega}$. The moments $P_{a_1 a_2 \dots a_n}(\Lambda)$ will now be encoded in the trigonometric polynomial given by the direction dependent limit $\lim_{\rho \rightarrow 0} f_n(\rho, \phi)$, which takes the form in (3). See also lemma 3. Note that although $P_{a_1 a_2 \dots a_n}$ is analytic on $\hat{V}$, f_n will not be analytic in terms of ξ, η since η^a is direction dependent at $\Lambda \in S$. In general we have the following lemmas:

Lemma 2. *Suppose that f is an analytic function, on a ball of radius r_0 around Λ on $\hat{V}$. Then the restriction of f to S , f_L , can be decomposed as $f_L(\xi, \zeta) = f_1(\xi, \zeta) + i\rho f_2(\xi, \zeta)$, where f_1 and f_2 are analytic in terms of ξ and ζ on the disk $\xi^2 + \zeta^2 < \frac{r_0^2}{2}$, and where $\rho = \sqrt{\xi^2 + \zeta^2}$. Furthermore, if f is real-valued then f_1 and f_2 are real-valued.*

Proof. We start by splitting $f = f(x, y, z)$ into its even, f_e , and odd, f_o , part with respect to z . We can now rewrite $f_e(x, y, z) = \tilde{f}_e(x, y, z^2)$ and $f_o(x, y, z) = z\tilde{f}_o(x, y, z^2)$, where both $\tilde{f}_e$ and $\tilde{f}_o$ are analytic in their arguments (at least near $(0, 0, 0)$). The restriction of $\tilde{f}_e$ to S gives f_1 : $f_1(\xi, \zeta) = \tilde{f}_e(\xi, \zeta, -(\xi^2 + \zeta^2))$, while the restriction of $\tilde{f}_o$ gives f_2 : $f_2(\xi, \zeta) = \tilde{f}_o(\xi, \zeta, -(\xi^2 + \zeta^2))$. Adding these, and also noting that $z \rightarrow i\rho$ gives the required decomposition. On S we have $|z|^2 = \xi^2 + \zeta^2$, hence $\xi^2 + \zeta^2 < \frac{r_0^2}{2}$ implies $x^2 + y^2 + z^2 < r_0^2$. This gives the domain of analyticity. The reality follows from the construction. $\square$

Remark: In f_L , the subscript L stands for the 'leading term'.

Although tensor fields on $\hat{V}$ can be pulled back to $S \setminus \{\Lambda\}$, we will only need their contractions with the appropriate number of η^a vectors. This contraction will introduce a direction dependence which shows up in the following lemma.

Lemma 3. Suppose $T_{a\dots b}$ is an analytic tensor field on a ball of radius r_0 around Λ on $\hat{V}$. Then the scalar field $f_L = \eta^a \dots \eta^b T_{a\dots b}$ on S can be written as $f_L(\xi, \zeta) = \frac{1}{\rho^n}(f_1(\xi, \zeta) + i\rho f_2(\xi, \zeta))$, where f_1 and f_2 are analytic (on the disk with radius $\frac{r_0}{\sqrt{2}}$ around the origin) in terms of ξ and ζ , and where $\rho = \sqrt{\xi^2 + \zeta^2}$. Furthermore, if $T_{a\dots b}$ is real-valued then f_1 and f_2 are real-valued.

Proof. Consider the scalar field $g = x^a \dots x^b T_{a\dots b}$ on $\hat{V}$, where

$$x^a = x \left(\frac{\partial}{\partial x} \right)^a + y \left(\frac{\partial}{\partial y} \right)^a + z \left(\frac{\partial}{\partial z} \right)^a.$$

This scalar field is analytic on the same ball as $T_{a\dots b}$. Hence we can use lemma 2 to obtain analytic functions f_1 and f_2 such that $g_L = f_1 + i\rho f_2$. The reality also follows from lemma 2. Furthermore, $x^a = \rho\eta^a$ on S . Thus $g_L = \rho^n f_L$. But f_L is bounded near the origin, thus we can divide g_L by ρ^n and get the lemma. $\square$

We remark that the boundedness of $f_L(\xi, \zeta) = \frac{1}{\rho^n}(f_1(\xi, \zeta) + i\rho f_2(\xi, \zeta))$ when $\rho \rightarrow 0$ implies that both f_1 and f_2 have zeros of sufficient order at $(\xi, \zeta) = (0, 0)$. It also implies that f_L will be direction dependent there.

We can now contract (1) with η^a and get the following theorem.

Theorem 4. Let $\hat{V}$ and S be defined as in sections (2) and (3). Let η^a have the properties given by lemma 1, and let f_n be defined by (4). Then the recursion (1) on $\hat{V}$ takes the form

$$f_n = \eta^a \hat{D}_a f_{n-1} - \frac{(n-1)(2n-3)}{2} \eta^a \eta^b \hat{R}_{ab} f_{n-2} \quad (5)$$

on S . The moments of order 2^n are captured in the direction dependent limit $\lim_{\rho \rightarrow 0} f_n(\rho, \varphi)$.

Proof. That (1) takes the form (5) follows exactly as in [4] using that $\eta^a \hat{D}_a \eta^b = 0$, although the recursion is defined only on S rather than on $\hat{V}$. The last statement is the content of lemma 1c. $\square$

3.1 Simplified calculation of the moments

In this section, we will show that it is possible to obtain the recursion (5) without the term involving the Ricci tensor. This will be accomplished by using the conformal freedom at hand, i.e. change Ω . The conformal freedom is $\Omega \rightarrow \tilde{\Omega} = \frac{\Omega}{\alpha}$ where α is analytic near Λ with $\alpha(\Lambda) = 1$ ². $\hat{D}_a(\frac{1}{\alpha})$ at Λ gives a shift of the moments which corresponds to a 'translation' of the physical space, [6]. Hence we can assume that $\hat{D}_a(\frac{1}{\gamma}) = 0$ at Λ . It is to be noted that a change of Ω changes the (rescaled) potential $\phi/\sqrt{\Omega}$. It also changes the normal coordinates on $\hat{V}$, and hence all conclusions must be made with some care. In order to derive the simplified recursion (14), we will specify a α through α_L , the restriction of α to S . However, in order to deduce that there exists a *real-valued* function α which the prescribed values of α_L , we need to say more on the representation of α_L . This result and a useful estimate is the content of lemma 5.

²A more natural condition is the equivalent statement $\Omega \rightarrow \tilde{\Omega} = \Omega\alpha$. However, this formulation gives slightly neater calculations.

Lemma 5. Let $f_L = f_1(\xi, \zeta) + i\rho f_2(\xi, \zeta)$ where f_1 and f_2 are analytic on the ball $U = \{|\xi|^2 + |\zeta|^2 < r_0^2\}$, and where $\rho = \sqrt{\xi^2 + \zeta^2}$. We can then write

$$f_L(\rho \cos \varphi, \rho \sin \varphi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l c_{l,m} e^{im\varphi} \rho^l, \quad (6)$$

where

$$\sum_{l=0}^{\infty} \sum_{m=-l}^l |c_{l,m}| \rho^l < \infty, \quad \rho < r_0. \quad (7)$$

Furthermore, the converse is true; if f_L is a function satisfying (6) and (7) then there are functions f_1 and f_2 analytic in U such that $f_L = f_1 + i\rho f_2$. The functions f_1 and f_2 are real-valued if and only if the coefficients $c_{l,m}$ satisfy $\bar{c}_{l,m} = (-1)^{l-m} c_{l,-m}$.

Proof. Form the functions $f_1(\frac{x+y}{2}, \frac{x-y}{2i})$ and $f_2(\frac{x+y}{2}, \frac{x-y}{2i})$.³ These functions are analytic in terms of x and y . Therefore there exist coefficients $c_{l,m}$ such that

$$f_j(\frac{x+y}{2}, \frac{x-y}{2i}) = \sum_{l=0}^{\infty} \sum_{k=0}^l i^{1-j} c_{l+j-1, 2k-l} x^k y^{l-k}, \quad (8)$$

where $j = 1$ or 2 , and

$$\sum_{l=0}^{\infty} \sum_{k=0}^l |c_{l+j-1, 2k-l}| |x|^k |y|^{l-k} < \infty, \quad |\frac{x+y}{2}|^2 + |\frac{x-y}{2i}|^2 = \frac{1}{2}(|x|^2 + |y|^2) < r_0^2.$$

Now let $x = \rho e^{-i\varphi}$ and $y = \rho e^{i\varphi}$. For all $\rho^2 = \frac{1}{2}(|x|^2 + |y|^2) < r_0^2$ this shows that

$$f_L(\rho \cos \varphi, \rho \sin \varphi) = \sum_{l=0}^{\infty} \sum_{k=0}^l c_{l, 2k-l} e^{i(2k-l)\varphi} \rho^l + \sum_{l=0}^{\infty} \sum_{k=0}^l c_{l+1, 2k-l} e^{i(2k-l)\varphi} \rho^{l+1}. \quad (9)$$

The reality condition follows easily from (8) and (9). A reorganisation gives the first part of the lemma. The converse is given by the substitution $x = \xi + i\zeta$ and $y = \xi - i\zeta$ in (8). The series converges absolutely for $|\xi|^2 + |\zeta|^2 = \rho^2 < r_0^2$. $\square$

We now show that we can choose α such that the Ricci term in (5) vanishes.

Lemma 6. There exists a real-valued real analytic function α on $\hat{V}$ such that the Ricci tensor $\tilde{R}_{ab}$ of the new metric $\tilde{h}_{ab} = \alpha^{-2} \hat{h}_{ab}$ satisfies $\tilde{\eta}^a \tilde{\eta}^b \tilde{R}_{ab} = 0$ on $\tilde{S}$, where both $\tilde{\eta}^a$, $\tilde{S}$ and the mapping F are defined in terms of coordinates which are normal with respect to the new metric $\tilde{h}_{ab}$.

Proof. We first demonstrate that we can make $\eta^a \eta^b \tilde{R}_{ab} = 0$ on S . From [10]

$$\tilde{R}_{ab} = \hat{R}_{ab} + \frac{1}{\alpha} \hat{D}_a \hat{D}_b \alpha + \alpha^{-1} \hat{h}_{ab} \hat{h}^{cd} \hat{D}_c \hat{D}_d \alpha - 2\alpha^{-2} \hat{h}_{ab} \hat{h}^{cd} (\hat{D}_c \alpha) \hat{D}_d \alpha.$$

On S , (using η^a and S belonging to $\hat{h}_{ab}$), this becomes

$$\eta^a \eta^b \tilde{R}_{ab} = \eta^a \eta^b \hat{R}_{ab} + \frac{1}{\alpha_L} \eta^a \eta^b \hat{D}_a \hat{D}_b \alpha_L = \eta^a \eta^b \hat{R}_{ab} + \frac{1}{\alpha_L} \frac{\partial^2 \alpha_L}{\partial \rho^2} \quad (10)$$

³Although we will always take ξ and ζ to be real, they are temporarily complexified in this proof.

where we have used $\eta^a \hat{D}_a \eta^b = 0$. In order to make $\eta^a \eta^b \tilde{R}_{ab} = 0$, we require

$$\alpha_{L\rho\rho} + \eta^a \eta^b \hat{R}_{ab} \alpha_L = 0. \quad (11)$$

Note that, on $\hat{V}$, $\hat{R}_{ab}$ satisfies the conditions of lemma 3, therefore from lemma 3 and lemma 5 it follows that

$$\eta^a \eta^b \hat{R}_{ab} = \sum_{n=0}^{\infty} b_{n+2} (e^{i\varphi}, e^{-i\varphi}) \rho^n, \quad (12)$$

where each b_n is a polynomial of degree at most n . It is easy to see that the reality condition from lemma 5 is equivalent to $\overline{b_n(e^{i\varphi}, e^{-i\varphi})} = (-1)^n b_n(-e^{i\varphi}, -e^{-i\varphi})$. Hence the reality of $\hat{R}_{ab}$ implies this condition. Moreover, since $\hat{R}_{ab}$ is analytic in a neighbourhood of Λ , there exists a ρ_0 such that for any fixed φ , the series (12) converges for all $\rho < \rho_0$. Well known ODE theory gives that for each fixed φ there exists a solution α_L , analytic in ρ for $0 \leq \rho < \rho_0$, i.e., for all φ we have

$$\alpha_L = \sum_{n=0}^{\infty} a_n(\varphi) \rho^n, \quad \rho < \rho_0.$$

We note that $a_0 = 1$ while a_1 can be chosen to be 0 (translation). α_L will have the right regularity if we can show that each $a_n(\varphi)$ is a polynomial in $e^{i\varphi}$ and $e^{-i\varphi}$ of degree at most n . Equation (11) becomes

$$-\sum_{n=0}^{\infty} n(n-1) a_n \rho^{n-2} = \sum_{n=0}^{\infty} \sum_{j=0}^{\infty} b_{n+2} a_j \rho^{n+j}.$$

Equating powers of ρ we get

$$-(n+2)(n+1) a_{n+2} = \sum_{m=0}^n b_{m+2} a_{n-m}, \quad n \geq 0. \quad (13)$$

The polynomials a_0 and a_1 has maximal degree 0 respective 1. The polynomials b_n has maximal degree n . Straightforward induction shows that a_n has maximal degree n . Induction and (13) also implies $\overline{a_n(e^{i\varphi}, e^{-i\varphi})} = (-1)^n a_n(-e^{i\varphi}, -e^{-i\varphi})$. Thus, due to lemma 5, there are real-valued analytic functions α_1 and α_2 such that $\alpha_L = \alpha_1 + i\rho\alpha_2$. The function $\alpha = \alpha_1(x, y) + z\alpha_2(x, y)$ is then a real-valued analytic extension of α_L to a ball in $\hat{V}$. This shows that there exist an α such that $\eta^a \eta^b \tilde{R}_{ab} = 0$ on S .

However, it then also follows that $\tilde{\eta}^a \tilde{\eta}^b \tilde{R}_{ab} = 0$ on $\tilde{S}$. From α we get $\tilde{\Omega} = \Omega/\alpha$, and the corresponding new metric $\tilde{h}_{ab} = \alpha^{-2} \hat{h}_{ab}$. Note that in (5), where now $\eta^a \eta^b \hat{R}_{ab} = 0$, the recursion is stated in terms of $\hat{D}_a$, i.e., it is expressed in terms of $\hat{h}_{ab}$ instead of $\tilde{h}_{ab}$. From $\tilde{h}_{ab}$ we get new normal coordinates, i.e., (x, y, z) in $T_\Lambda \hat{V}_\mathbb{C}$, are mapped into $\hat{V}_\mathbb{C}$ using the exponential map belonging to $\tilde{h}_{ab}$. We then construct $\tilde{\eta}^a$ and the mapping F with respect to these coordinates. Now, null geodesics, of which S consists, are conformally invariant, although they become non-affinely parametrised. This means that $\tilde{\eta}^a \propto \eta^a$, and that points in S are mapped into points in $\tilde{S}$. Thus $\tilde{\eta}^a \tilde{\eta}^b \tilde{R}_{ab} \propto \eta^a \eta^b \tilde{R}_{ab} = 0$ on $\tilde{S}$ (or S). $\square$

Henceforth we denote all entities defined via $\tilde{h}_{ab}$ instead of $\hat{h}_{ab}$ with a tilde. In particular, $\tilde{D}_a$ will denote the derivative operator associated with $\tilde{h}_{ab}$. Applying lemma 6 to theorem 4 we immediately get the following theorem.

Theorem 7. *Let $\hat{V}$ and $\tilde{S}$ be defined as in sections (2) and (3), where $\tilde{S}$ is defined in terms of normal coordinates connected to $\tilde{h}_{ab}$. Let $\tilde{\eta}^a$ have the properties given by lemma 1 with respect to $\tilde{h}_{ab}$, and let $\tilde{f}_n$ be defined by (4) with $\tilde{\eta}^a$ replacing η^a . Then the recursion (1) on $\hat{V}$ takes the form*

$$\tilde{f}_n = \tilde{\eta}^a \tilde{D}_a \tilde{f}_{n-1} = (\tilde{\eta}^a \tilde{D}_a)^n \tilde{f}_0 = \frac{\partial^n}{\partial \tilde{\rho}^n} \tilde{f}_0 \quad (14)$$

on $\tilde{S}$. The moments of order 2^n are captured in the direction dependent limit $\lim_{\tilde{\rho} \rightarrow 0} \tilde{f}_n(\tilde{\rho}, \varphi)$.

Proof. Since $\tilde{\eta}^a \tilde{D}_a$ is $\frac{\partial}{\partial \tilde{\rho}}$, each $\tilde{f}_n$ is easily derived from $\tilde{f}_0$. Also, from lemma 1c), we (again) know that the $2n + 1$ degrees of freedom of $P_{a_1 \dots a_n}$ at Λ are encoded in $\tilde{f}_n$. $\square$

4 Bounds on the moments

All multipole moments are encoded in $\tilde{f}_0$, and we note that the recursion (14) is identical to the recursion emanating from a scalar function in $\mathbb{R}^3$ (after inversion). It is clear that many different functions on $\mathbb{R}^3$ will produce the same moments, but if we also require that the function, g say, is harmonic, g is uniquely determined by the moments.

Thus, provided that we can connect a function which is harmonic in a neighbourhood of $\mathbf{0} \in \mathbb{R}^3$ to each $\tilde{f}_0$, we have the following theorem:

Theorem 8. *Suppose that (M, g_{ab}) is a stationary asymptotically flat space-time, admitting an analytic (rescaled) potential and an analytic chart⁴ on the conformally compactified manifold of timelike Killing trajectories, around the infinity point Λ . Then there exist a (flat-)harmonic function g in a neighbourhood of $\mathbf{0} \in T_\Lambda \hat{V} \cong \mathbb{R}^3$, such that all multipole moments of M are given by*

$$P_{a_1 \dots a_n}(\Lambda) = (\nabla_{a_1} \dots \nabla_{a_n} g)(0), \quad (15)$$

where ∇_a in the LHS of (15) is the flat derivative operator in $\mathbb{R}^3$. This puts a bound on the multipole moments, since the Taylor expansion $\sum_{|\alpha| \geq 0} \frac{\mathbf{r}^\alpha}{\alpha!} (\partial_\alpha g)(\mathbf{0})$ of g converges in a neighbourhood of the origin in $\mathbb{R}^3$.

Proof. By lemma 5 we have

$$\tilde{f}_L(\rho \cos \varphi, \rho \sin \varphi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l c_{l,m} e^{im\varphi} \rho^l, \quad \rho < r_0$$

for some $r_0 > 0$, and we know that $\tilde{f}_L$ fully determines the moments of M . We will now define the function g , which will be shown to be harmonic in a neighbourhood of the origin. Finally we will argue that g has the correct

⁴As discussed before, the analyticity has been proved for the case with non-zero mass.

derivatives at $\mathbf{0}$, i.e., that the equality (15) is valid (for all $n \geq 0$). First we define the coefficients

$$a_{l,m} = c_{l,m} i^{-m-l} 2^{1-l} \pi \frac{\sqrt{(l+m)!(l-m)!}}{\Gamma(l + \frac{1}{2}) \sqrt{2l+1}}.$$

and the function

$$g(r, \theta, \varphi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l a_{l,m} Y_l^m(\theta, \varphi) r^l.$$

Due to the construction, g is harmonic at those (interior) points for which the sum converges. We now establish convergence. From the identity

$$\sum_{m=-l}^l |Y_l^m(\theta, \varphi)|^2 = \frac{2l+1}{4\pi}$$

we get

$$\begin{aligned} \left| \sum_{m=-l}^l a_{l,m} Y_l^m \right| &\leq \sum_{m=-l}^l |a_{l,m} Y_l^m| \leq \left(\sum_{m=-l}^l |a_{l,m}|^2 \right)^{\frac{1}{2}} \left(\sum_{m=-l}^l |Y_l^m|^2 \right)^{\frac{1}{2}} \\ &= \left(\sum_{m=-l}^l \left| \sqrt{\frac{2l+1}{4\pi}} a_{l,m} \right|^2 \right)^{\frac{1}{2}} = \frac{\sqrt{\pi}}{2^l \Gamma(l + \frac{1}{2})} \left(\sum_{m=-l}^l \left| c_{l,m} \sqrt{(l+m)!(l-m)!} \right|^2 \right)^{\frac{1}{2}} \end{aligned}$$

Furthermore, the inequality $(l+m)!(l-m)! \leq (2l)!$, when $-l \leq m \leq l$ follows from the convexity of $\ln(\Gamma(x))$. Therefore

$$\left| \sum_{m=-l}^l a_{l,m} Y_l^m \right| \leq \frac{\sqrt{\pi} \sqrt{(2l)!}}{2^l \Gamma(l + \frac{1}{2})} \left(\sum_{m=-l}^l |c_{l,m}|^2 \right)^{\frac{1}{2}}$$

Next, the inequality $\frac{\pi(2l)!}{4^l \Gamma(l + \frac{1}{2})^2} = \frac{(2l)!!}{(2l-1)!!} = (2l+1) \frac{(2l)!!}{(2l+1)!!} \leq 2l+1$ gives

$$\left| \sum_{m=-l}^l a_{l,m} Y_l^m \right| \leq \sqrt{2l+1} \left(\sum_{m=-l}^l |c_{l,m}|^2 \right)^{\frac{1}{2}} \leq \sqrt{2l+1} \sum_{m=-l}^l |c_{l,m}|$$

But for all $\epsilon > 0$ we have $\sqrt{2l+1}(1+\epsilon)^{-l} \rightarrow 0$ as $l \rightarrow \infty$. So the factor $\sqrt{2l+1}$ will not affect the radius of convergence.

Hence

$$\sum_{l=0}^{\infty} \sum_{m=-l}^l a_{l,m} Y_l^m r^l \text{ converges if } r < r_0.$$

This shows that g is well defined in a neighbourhood of $\mathbf{0}$ in $T_{\Lambda} \hat{V}$, and we must now show that we have equality in (15). We will do this by forming g_L and then compare with $\tilde{f}_L$. Note, however, that $\tilde{f}_L$ is defined on $\tilde{S} \subset V_{\mathbb{C}}$, while g_L will be defined on the corresponding surface $\tilde{S} = S_{\tilde{h}_{ab}(\Lambda)}$ in $T_{\Lambda} \hat{V}_{\mathbb{C}}$. (By $\tilde{S}$ we denote the surface defined by F as previously, but where F now maps (ξ, ζ) into $T_{\Lambda} \hat{V}_{\mathbb{C}}$ via the flat metric $\tilde{h}_{ab}(\Lambda)$.) This means that f_L and g_L really can be

compared only at Λ (i.e. $\mathbf{0} \in T_\Lambda \hat{V}$), where also the equality (15) is evaluated. On the other hand, the radial derivatives of both entities are well defined and comparable at Λ . In other words, if both $\tilde{f}_L$ and g_L are equal when expressed in terms of the coordinates ξ, ζ , they will produce the same derivatives/moments. To summarise; in the case of g , the function F and the vector η^a are simply interpreted in $T_\Lambda \hat{V}_\mathbb{C}$ rather than in $V_\mathbb{C}$.

To proceed, we recall that

$$Y_l^m(\theta, \varphi) = i^{m-|m|} \sqrt{\frac{(2l+1)(l-|m|)!}{4\pi(l+|m|)!}} P_l^{|m|}(\cos \theta) e^{im\varphi}$$

for $-l \leq m \leq l$, and that

$$P_l^m(\cos \theta) = (-1)^m 2^{-l} \sin^m \theta \sum_{k=0}^{\lfloor \frac{l-m}{2} \rfloor} \frac{(-1)^k (2l-2k)!}{k!(l-k)!(l-2k-m)!} \cos^{l-m-2k} \theta$$

where $0 \leq \theta \leq \pi$ and $m \geq 0$. Therefore,

$$g(r, \theta, \varphi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l \sum_{k=0}^{\lfloor \frac{l-|m|}{2} \rfloor} \frac{c_{l,m} \sqrt{\pi} (l-|m|)! (2l-2k)! e^{im\varphi} \rho^{|m|} z^{l-|m|-2k} r^{2k}}{i^{l-|m|} (-1)^k 4^l k! (l-k)! (l-|m|-2k)! \Gamma(l + \frac{1}{2})},$$

where $z = r \cos \theta$, $\rho = r \sin \theta$. When we take the restriction of g to $\bar{S}$, i.e., form g_L , only the terms with $k = 0$ survives since $r_L = 0$. Thus

$$g_L(\rho \cos \theta, \rho \sin \theta) = \sum_{l=0}^{\infty} \sum_{m=-l}^l \frac{c_{l,m} \sqrt{\pi} (2l)! e^{im\varphi} \rho^l}{4^l l! \Gamma(l + \frac{1}{2})} = \sum_{l=0}^{\infty} \sum_{m=-l}^l c_{l,m} e^{im\varphi} \rho^l = \tilde{f}_L,$$

which means that we have equality in (15). $\square$

5 Discussion

In this paper we have studied the multipole moments of stationary asymptotically flat spacetimes. By using normal coordinates, and by exploiting the conformal freedom, we could show that the tensorial recursion (1) could be replaced by the scalar recursion (14). This recursion is a direction dependent recursion on $\mathbb{R}^2$, where the moments are encoded in the direction dependent limits at Λ .

Using this setup, we could also show that the multipole moments cannot grow too fast. In essence, the rescaled potential behaves (locally) in the manner of a harmonic function on $\mathbb{R}^3$. The bounds on the moments given in theorem 8 gives the necessary part in a conjecture due to Geroch [6], and it is of course tempting to conjecture that this condition on the moments also will be sufficient (as long as the monopole is real-valued).

Whether this can be proved using the techniques presented here is still an open question.

We also remark that similar questions concerning the convergence of asymptotic expansions in the static case are currently being studied by Friedrich, using a different technique, [5].

References

- [1] Beig, R., *The Multipole Expansion in General Relativity*, Acta Physica Austriaca, **53**, 249 (1981).
- [2] Beig, R., Simon, W., *On the multipole expansion for stationary spacetimes*, Proc. R. Soc. Lond. A, **376**, 333 (1981).
- [3] Bäckdahl, T., Herberthson, M., *Static axisymmetric spacetimes with prescribed multipole moments*, Class. Quantum Grav. **22**, 1607 (2005).
- [4] Bäckdahl, T., Herberthson, M., *Explicit multipole moments of stationary axisymmetric spacetimes*, Class. Quantum Grav. **22**, 3585 (2005).
- [5] Friedrich, H., *On the convergence of certain expansions at space-like infinity of asymptotically flat, static vacuum solutions*, <http://www.newton.cam.ac.uk/webseminars/pg+ws/2005/gmr/1202/friedrich/>
- [6] Geroch, R., *Multipole Moments. II. Curved Space*, J. Math. Phys., **11**, 2580 (1970).
- [7] Hansen, R.O., *Multipole moments of stationary spacetimes*, J. Math. Phys., **15**, 46 (1974).
- [8] Herberthson, M., *The gravitational dipole and explicit multipole moments of static axisymmetric spacetimes*, Class. Quantum Grav. **21**, 5121 (2004).
- [9] Kundu, P., *On the analyticity of stationary gravitational fields at spatial infinity*, J. Math. Phys., **22**, 2006 (1981).
- [10] Wald, R.M., *General Relativity*, The University of Chicago Press, Chicago 1984.